Protecting Your Bones During Menopause

Osteoporosis Prevention, Diagnosis, and Treatment

What to expect—and what to accept?

A simple question-and-answer guide for women

Menopause is a natural stage of life, not an illness. Every woman's experience is different. Some women have few symptoms, while others find the changes seriously interfere with work and home life.

This leaflet summarises information from the published medical documents. It is intended for education and discussion with a healthcare professional. It does not replace personal medical advice.

Menopause is a significant life transition that brings about various changes, some more widely discussed than others. While hot flashes, mood swings, and sleep disturbances often take centre stage, there's a less obvious but equally critical concern that every woman should be aware of: her bone health. The rapid decline in oestrogen during menopause can profoundly impact your skeletal system, increasing your risk of developing osteopenia and, eventually, osteoporosis.

Osteoporosis is often called the "silent thief" because it can progress without noticeable symptoms until a fracture occurs, dramatically impacting your quality of life and independence. But the good news is, understanding this evolving landscape of bone health in midlife means you can take proactive steps for prevention, accurate diagnosis, and effective management.

This article, presented in an easy-to-understand question-and-answer format, will provide you with essential scientific facts and practical advice to safeguard your bones during and after menopause.

Q: What exactly is menopause, and why does it matter for my bones?

Menopause marks the *end of a woman's reproductive years*, officially diagnosed after 12 consecutive months without a menstrual period. It typically occurs ***between the ages of 45 and 55***, with the average age being 46.2. This transition is characterized by *a significant decrease in the hormone oestrogen*, primarily produced by the ovaries.

Oestrogen plays a crucial role in maintaining strong bones. It helps inhibit osteoclasts, the cells responsible for breaking down old bone tissue. When estrogen levels drop sharply during menopause, this natural "brake" on bone breakdown is released. This leads to ***bone resorption (breakdown) happening faster than bone formation, resulting in a rapid loss of bone mineral density (BMD).*** Research indicates that ***women can lose anywhere from 10% to 20% of their total bone density within the first five to seven years*** after menopause begins. This rapid decline makes midlife a critical period for bone health intervention.

Q: I had early menopause (before age 45). Am I at higher risk?

Yes, absolutely. If you experience menopause early, whether naturally, due to primary ovarian insufficiency, or surgical intervention (e.g., hysterectomy with ovary removal), your bones are exposed to estrogen deficiency for a longer period. This **extended deficit accelerates the rate of bone loss**, often leading to lower peak bone mass and a heightened risk of developing osteoporosis at a younger age. Proactive monitoring and discussions with your doctor are especially important in this situation.

Q: My doctor mentioned "bone remodelling." What is that, and what are osteopenia and osteoporosis?

A: Your **bones are living tissues constantly undergoing a process called remodelling**. Old bone tissue is continuously broken down by cells called osteoclasts, and new bone tissue is formed by cells called osteoblasts. This cycle helps keep your bones strong and repair minor damage. In childhood and early adulthood, bone formation outpaces breakdown, leading to **peak bone mass (the strongest your bones will ever be), typically reached between ages 25-30**. After age 40, bone mass slowly decreases.

During menopause, the drop in oestrogen causes the breakdown process to accelerate beyond the body's ability to rebuild, leading to **a net loss of bone.**

- **Osteopenia:** This is a condition where your **bone mineral density is lower than the average peak** but not yet severe enough to be classified as osteoporosis. Think of it as a ***warning sign*** – your bones are weakening, and it's time for intervention to prevent it from worsening.
- **Osteoporosis:** Meaning ***"porous bone,"*** this is a systemic skeletal disorder where bones become **very weak, brittle, and structurally compromised**. The internal honeycomb-like structure of the bone develops larger spaces and gaps. **This significantly increases the risk of fractures, even from minor falls or everyday activities**. Osteoporosis is often described as a **"silent disease"** because it **usually has no symptoms until a bone breaks**. Over time, it can lead to physical changes like **loss of height** or **a stooping posture** due to spinal compression fractures.

Q: How common are fractures due to osteoporosis in menopausal women?

A: The statistics are sobering: approximately one in two women over the age of 60 **will experience at least one fracture due to osteoporosis**. These fractures, particularly in the hip or spine, can lead to severe pain, decreased mobility, loss of independence, and a reduced quality of life. *Preventing them is a primary goal of osteoporosis management*.

Q: How will my doctor assess my bone health and diagnose osteoporosis?

A: The gold standard for measuring bone mineral density (BMD) and diagnosing osteoporosis is a **DXA (Dual-energy X-ray Absorptiometry) scan**. This is a **quick, non-invasive imaging test** that uses **very low doses of radiation** to measure the density of your bones, typically in the lower spine and hip.

The DXA scan provides a **T-score**, which compares your BMD to that of a healthy young adult (around 30-35 years old).

- A T-score of **-1.0 or above** is considered normal.
- A T-score between **-1.0 and -2.5** indicates osteopenia.
- A T-score of **-2.5 or lower** confirms a diagnosis of osteoporosis.

Q: When should I get a DXA scan?

A: General recommendations suggest:

- All women aged **65 years and older** should undergo a bone density study.
- Women aged **60 to 64 with increased fracture risk** (e.g., a history of fractures or a family history of osteoporosis) should also be screened.
- If you have **specific risk factors**, such as early menopause (before 45), certain medical conditions, or medications like long-term corticosteroids, your doctor may recommend earlier screening, even if you are younger.

Your doctor may also use a **FRAX (Fracture Risk Assessment) tool** to **calculate your 10-year probability of experiencing a major osteoporotic fracture**, considering various risk factors beyond just your T-score. Additionally, **blood tests** for Vitamin D, calcium, thyroid function, and other markers may be conducted to get a comprehensive picture of your bone health.

Q: What can I do in my daily life to prevent bone loss and keep my bones strong?

A: **Lifestyle changes** are foundational for bone health, especially during and after menopause. Even if you need medication, these habits significantly support your skeletal system:

- **Nutrition is Key:**
    - **Calcium:** Aim for **1,200 milligrams of calcium daily** if you're over 50. Excellent sources include dairy products (milk, yogurt, cheese), firm tofu, almonds, dark green leafy vegetables (like kale and spinach), and fish with edible bones (like sardines).
    - **Vitamin D:** This **"sunshine vitamin"** is crucial for your body to absorb calcium effectively. The recommendation is **600-800 International Units (IU) per day**, increasing to **800 IU for those over 70.** While sunlight exposure helps, many people, especially in regions with limited sun or with darker skin, may need supplements. Dietary sources include **oily fish (salmon, mackerel) and egg yolks**, as well as **fortified foods.**
    - **Balanced Diet:** Include **lean proteins, whole grains, fruits, and vegetables** to provide other vital nutrients like magnesium, vitamin K, and vitamin C, all important for bone health. ***A Mediterranean-style diet can be very beneficial.***
- **Stay Active:** Exercise puts beneficial stress on your bones, stimulating bone remodelling and making them stronger.
    - **Weight-Bearing Exercise:** Activities where you're on your feet and working against gravity, such as **brisk walking, dancing, jogging, hiking, or climbing stairs**.
    - **Resistance Training:** Using weights, resistance bands, or your own body weight (e.g., squats, push-ups) builds muscle, which in turn strengthens bones. Aim for ***2-3 sessions per week*** on non-consecutive days.
    - **Balance and Flexibility:** Practices like **yoga, Tai Chi, and Pilates** improve coordination and balance, **significantly reducing the risk of falls,** which are a major cause of fractures.
    - **Consult a Professional:** Before starting any new exercise regimen, especially if you have existing bone density loss, consult your doctor or a physical therapist to ensure it's safe and effective for you.
- **Avoid Harmful Habits:**
    - **Smoking:** Smoking significantly slows down bone-building cells and increases fracture risk. Quitting can help reverse this risk.
    - **Alcohol:** Excessive alcohol intake can weaken bones and increase the risk of falls. Limit consumption to recommended guidelines (no more than 14 units per week, with regular alcohol-free days).
    - **Caffeine:** High caffeine intake may also contribute to bone loss, so moderation is advised.

Q: If lifestyle changes aren't enough, what medical treatments are available for osteoporosis?

A: There's an evolving landscape of effective medications designed to slow bone loss, **increase bone density**, and **reduce fracture risk.** Your doctor will work with you to determine the best approach based on your individual needs and risk profile.

- **Hormone Replacement Therapy (HRT) / Menopausal Hormone Therapy (MHT):**
    - **How it works:** By replacing the oestrogen your body is no longer producing, HRT can effectively prevent bone loss and help maintain bone density. It also significantly alleviates other menopausal symptoms like hot flashes and vaginal dryness.
    - **When it's used:** HRT is often a beneficial option when started around the time of menopause, especially for women with bothersome menopausal symptoms who also need bone protection. Recent studies show that ***early HRT can significantly reduce the risk of osteoporosis and fractures.***
    - **Considerations:** HRT is not suitable for everyone (e.g., women with certain types of cancer). **Risks versus benefits,** including cardiovascular health and breast cancer risk, should be thoroughly discussed with your doctor. Transdermal estrogen may be considered for women with a risk of blood clots.
- **Bisphosphonates (e.g., Alendronate, Risedronate, Zoledronic Acid, Ibandronate):**
    - **How they work:** These are **often the first-line treatment**. They work by binding to bone minerals and inhibiting the activity of osteoclasts, thus slowing down bone breakdown and allowing for a net gain in bone mass.
    - **Administration:** Available as oral pills (taken weekly or monthly) or intravenous (IV) infusions (given yearly or biannually).
    - **Considerations:** Generally effective at reducing fracture risk. Side effects can include gastrointestinal upset (for oral forms) or flu-like symptoms (for IV forms). Rare but **serious side effects** like **osteonecrosis of the jaw (ONJ) or atypical femur fractures (AFFs)** can occur, especially with long-term use. Your doctor might recommend a **"drug holiday" after 3-5 years** for some patients.
- **Denosumab (Prolia):**
    - **How it works:** A biological medication (monoclonal antibody) that **blocks a protein called RANKL, essential for osteoclast formation and function**. This significantly reduces bone resorption.
    - **Administration:** Given as a **subcutaneous (under the skin) injection every six months** by a healthcare professional.
    - **Considerations:** Highly effective, including for non-vertebral fractures. It's crucial to take calcium and vitamin D supplements, as it can cause low calcium levels. Un-

like bisphosphonates, it **cannot be stopped abruptly** without follow-up treatment, **as this can lead to rapid bone loss and multiple spinal fractures.**

- **Selective Oestrogen Receptor Modulators (SERMs) (e.g., Raloxifene):**
    - **How they work:** These medications ***act like estrogen in some body tissues (like bone, where they reduce bone resorption)*** but block estrogen's effects in others (like breast and uterine tissue).
    - **Considerations: Primarily reduces vertebral fracture risk** and may lower the risk of invasive breast cancer. Side effects can include hot flashes, leg cramps, and a small increased risk of blood clots.
- **Anabolic (Bone-Building) Agents (e.g., Teriparatide, Abaloparatide, Romosozumab):**
    - **How they work:** These are unique because they **actively stimulate new bone formation, rather than just slowing breakdown.**
    - **Administration:** Teriparatide and Abaloparatide are daily self-injections, typically **used for no more than two years** (due to a rare theoretical risk of bone cancer, observed in animal studies). After this period, an anti-resorptive medication is usually prescribed to maintain the newly built bone.
    - **Romosozumab (Evenity):** A newer option that acts by inhibiting sclerostin, a protein that interferes with bone formation. It's given ***as monthly injections for 12 months,*** followed by another bone medication. It's approved for high-risk postmenopausal women but carries a boxed warning about potential increased risk of heart attack or stroke.
    - **Considerations:** Generally reserved for individuals with severe osteoporosis, very low bone density, or those who have had multiple fractures or failed other treatments.

Q: My bone health seems complex. How do I ensure I get the right care?

A: The "one-size-fits-all" approach to bone health is outdated. Modern bone health management emphasizes individualized treatment plans and patient counselling. Here's how you can actively participate:

1. **Be Proactive:** Don't wait for a fracture to occur. Start discussing bone health with your primary care provider or gynaecologist as you approach or enter menopause. Early detection is key.
2. **Understand Your Risk Factors:** Discuss your personal and family medical history, lifestyle, and any medications you take. These factors are crucial for your doctor to assess your specific fracture risk.

3. **Engage in Shared Decision-Making:** Your treatment plan should be a collaboration between you and your healthcare team. Ask questions, understand the benefits and potential risks of all options (lifestyle changes, medications), and express your preferences.

4. **Regular Monitoring:** Adhere to recommended schedules for DXA scans and blood tests to track your bone density and overall bone health.

5. **Comprehensive Approach:** Recognize that bone health involves more than just medication. Combine medical treatments with consistent lifestyle modifications (diet, exercise, fall prevention) for the best outcomes.

Q: What are some questions I should ask my doctor about my bone health?

A: Here are a few important questions you might consider:

- "Based on my personal history and risk factors, when should I have my first DXA scan, or how often should I have one if I've already had one?"
- "What are my specific T-scores, and what do they mean for my fracture risk?"
- "Considering my other health conditions and preferences, what are the best osteoporosis prevention or treatment options for me?"
- "If I start a medication, how will we monitor its effectiveness, and what are the potential long-term side effects I should be aware of?"
- "Are there any specific exercises or dietary changes you recommend for my situation?"

Primary keywords: Menopause, Osteoporosis, Osteopenia

Secondary keywords: bone health, oestrogen decline, DXA scan, osteoporosis prevention, osteoporosis treatment, calcium and vitamin D, Hormone Replacement Therapy (HRT)

Key Takeaways: Why Early Action and Ongoing Management Matter

The journey through menopause is a critical time for bone health. The rapid **estrogen decline** can quickly lead to weakened bones, often silently. Understanding your risk, getting timely **DXA scans** for diagnosis, and implementing a comprehensive plan for **osteoporosis prevention** and **osteoporosis treatment** are paramount.

Remember, maintaining **bone health** is a lifelong endeavour, but midlife demands particular attention. With personalized guidance from your healthcare team, including appropriate **calcium and vitamin D** intake, targeted exercise, and potentially **Hormone Replacement Therapy (HRT)** or other medications, you can significantly reduce your risk of fractures. Early action and ongoing management are your best defence against osteoporosis, helping you maintain an active, independent, and vibrant life well beyond menopause.

Important: This leaflet summarises information from the published medical documents and does not replace individualised clinical assessment or the full guideline texts. It is intended for educational purpose, only. Because medical information is undergoing constant scrutiny and changing at a rapid pace, the readers are advised to check any information, that is current.

Compiled & Edited by: Team from Kolkata Menopause Society

1. **Dr Arnab Kr Basak**, MD., DNB., MNAMS., MRCOG.
2. **Dr Mandira Dasgupta,** MD., DNB., MNAMS., MRCOG., FMAS.
3. **Dr Ranjana Tibrewal,** MBBS(Hons), MS, (FIAOG , Diploma in Aesthetic Functional and Regenerative Gynaecology)
4. **Dr Shabana Roze Chowdhury,** MBBS., DMCW. Ph.D., FRSPH.
5. **Dr Somjita Chattopadhyay (Chakraborty),** MBBS., DGO.,. MS., DNB
6. **Dr Sukanta Misra,** MD.
7. **Dr Susmita Chattopadhyay,** MD,DNB,FRCOG,CCT(UK).